



## King's Research Portal

DOI:

[10.1038/s41415-019-0196-8](https://doi.org/10.1038/s41415-019-0196-8)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Coxon, J. D., Hosey, M-T., & Newton, J. T. (2019). The oral health of dentally phobic 12- and 15-year-olds: a descriptive analysis of the 2013 Child Dental Health Survey. *British Dental Journal*, 226(8), 595-599. <https://doi.org/10.1038/s41415-019-0196-8>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# **The oral health of dentally phobic 12 and 15 year olds: A descriptive analysis of the 2013 Child Dental Health Survey.**

Coxon J.D., Hosey M.T., Newton J.T

## **Key Points**

- Dentally phobic adolescent children are more likely to have dental decay.
- The oral health of children with a dental phobia is more likely to impact on their everyday life compared to non dental phobic children.
- Dentally phobic children are less likely to show behaviour related to the prevention of dental disease, such as attending the dentist for regular examinations or brushing twice a day.

## **Abstract**

**Introduction.** Dental Anxiety has been shown to be related to poorer oral health. Fewer data exist exploring the relationship between oral health status and dental anxiety in non-clinical populations in children.

**Aim.** To compare the oral health of phobic and non-phobic children aged 12-15 years.

**Design.** Secondary analysis of data from 12 year old and 15 year old children in the Child Dental Health Survey 2013.

**Setting.** National Survey

**Materials and Methods.** Participants were grouped into non phobic and phobic groups, depending on their self reported dental anxiety (MDAS). Descriptive analyses compared the two groups on social demographic factors, clinical status, self-reported oral health status, oral health related behaviours and oral health impact.

**Results** 601 children were classed as dentally phobic with 4144 classed as non phobic. Dental phobic children were more likely to be female, had more active decay and untreated dental disease, and rated their dental health as poorer. Phobic children were more likely to report that their oral health had a negative effect on their everyday life. This group were less likely to brush their teeth regularly or attend the dentist for check ups.

**Conclusions.** Dentally phobic children have more dental disease and express greater impact on their everyday life.

## **Introduction**

A fear of dental treatment is well documented in both the adult and child population <sup>(1,2)</sup> Dental fear is defined as a loss of control and anticipation that something traumatic will happen in relation to dentistry (3). Dental phobia is defined as a severe and out of proportion fear of dentistry leading to avoidance and impacting on everyday life.<sup>(4)</sup>

Previous research has suggested that dental fear in adults impacts on their daily life, <sup>(5,6,7)</sup> leading to psychological distress and even increased time on sick leave.<sup>(8,9)</sup> The research into children's dental anxiety and its effect on the child's oral health and oral health related behaviour is sparser.

The reported prevalence of dental anxiety in children differs depending on the study from 9% to 22% <sup>(3,10)</sup> Qualitative studies to examine what the child perceives when faced with the dental environment have been undertaken, <sup>(11)</sup> suggesting that a cognitive behavioural therapy assessment model may be useful to understand childhood dental anxiety and gained some insight in to what children perceive as aversive when attending the dentist. There have also been small-scale studies that suggest that the child's oral health related quality of life is worse if the child is dentally anxious.  
<sup>(12)</sup>

In the UK, the 2013 Child Dental Health Survey (CDHS) resulted in the publication of a broad overview on dental anxiety prevalence and relationships. Participants aged 12 years and 15 years old were asked to self-rate their anxiety levels and reported that 62% of 12 year olds and 54% of 15 year olds stated they had anxiety when attending the dentist; a further 14% of 12 year olds and 10% of 15 year olds stated they had extreme anxiety <sup>(13)</sup> There was no reported relationship between eligibility to free school meals, a marker of social deprivation but there was a difference between the genders, with girls more likely to report extreme anxiety.

The aim of this paper is to use the data generated by the highly refined methodology of the CDHS to perform a descriptive analysis of relationships between dental anxiety and gender, social deprivation, oral health status, oral health related behaviours and oral health related quality of life.

## **Materials and methods**

The study constituted an analysis of the data that was gathered during the 2013 CDHS. This was the fifth in a series of surveys, completed every 10 years, to look at the dental health of children in England, Wales and Northern Ireland. Questionnaires also allowed an insight into the children's experience of dentistry and their oral health related behaviours and attitudes. Further details on the methodology of the Child Dental Survey 2013 can be found at <https://files.digital.nhs.uk/publicationimport/pub17xxx/pub17137/cdhs2013-report1-attitudes-and-behaviours.pdf>

### *Assessment of fear.*

Participants aged 12 and 15 years were grouped into non-phobic and phobic groups depending on their self-reported dental anxiety. This was scored using the Modified Dental Anxiety Scale (MDAS). <sup>(14)</sup> This scale asked the participants to score how they felt in different scenarios, such as having a local anaesthetic, ranging from not anxious to extremely anxious on a five-point scale. Scores were summed and those participants with a total of 19 or above were ranked as phobic (n= 601) based on the work and validation of numerous studies.<sup>(15,16,17)</sup> Scores below were ranked as non-phobic (n = 4144).

### *Data analysis*

Social and demographic variables, variables found at clinical examination, the child's self-reported oral health status, behaviours that impact on oral health and oral health impact variables were tabulated (table 1) and grouped according to whether the children were phobic or not and analysed using SPSS (version 25). A simple statistical analysis was conducted using cross tabulation and chi-

squared test. As multiple hypotheses were being tested, a Bonferroni correction was applied to allow for the increased chance of a rare event leading to incorrectly rejecting the null hypothesis. Following this correction the result was deemed significant if  $P < 0.002$

Variables chosen were;

### **Social and demographic**

- Gender
- Free school dinner eligibility. This was used by the CDHS as a measure of poor socio-economic status. Parents can claim free school dinners for their children if they claim unemployment benefits, an income related support allowance or help due to immigration status.

### **Clinical findings**

- Number of decayed, missing, filled teeth (DMFT or decay experience), according to the 2003 criteria which states "All teeth with cavitated or visual dentine caries, restorations with cavitated or visual dentine caries, teeth with filled decay (otherwise sound) and teeth extracted due to caries. Excludes teeth with enamel caries present. The term obvious decay experience relates to teeth with dentinal cavities, missing teeth and filled teeth in the DMFT dental decay index." <sup>(18)</sup> This was grouped into two groups; no decay experience and decay experience.
- Active decay present in the mouth. This included caries with cavitation and caries with no cavitation. This was grouped into two groups of no decay present and decay present.
- Indications of soft tissue lesions: ulceration, fistula or abscess (PUFA). This was used as a measure of the clinical consequences of untreated dental caries, with the results grouped into a) no soft tissue lesion seen and b) soft tissue lesion seen.

### **Variables related to the child's own reported oral health status.**

These measures were gathered via a questionnaire given to the child.

- Child's self-rated dental health, scored on a scale of one to five rating from very good to very poor. The variable was divided into two groups; very good/good and fair/worse
- Child's self-rated general health, scored on a scale of one to five rating from very good to very poor. The variable was divided into two groups; "very good/good" and "fair/worse"
- Child reported toothache in the last 3 months, scored "yes" or "no" and grouped accordingly.
- Child reported sensitive tooth in the last 3 months, scored "yes" or "no" and grouped accordingly.
- Child reported bleeding or swollen gums in the last 3 months, scored "yes" or "no" and grouped accordingly.

- Child reported broken tooth in the last 3 months, scored “yes” or “no” and grouped accordingly.
- Child reported mouth ulcers in the last 3 months, scored “yes” or “no” and grouped accordingly.
- Child reported bad breath in the last 3 months, scored “yes” or “no” and grouped accordingly.

#### **Variables related to oral health related quality of life**

- CHILD-OIDP, an oral health-related quality of life index for children (19). This was broken down into eight aspects;
  - o Child reports difficulty eating in last 3 months
  - o Child reports difficulty speaking in last 3 months
  - o Child reports difficulty cleaning teeth in last 3 months.
  - o Child reports difficulty relaxing or sleeping in last 3 months
  - o Child reports that they felt different in last 3 months
  - o Child reports difficulty smiling, laughing, showing teeth in last 3 months.
  - o Child reports difficulty doing schoolwork in last 3 months.
  - o Child reports difficulty enjoying being with people in last 3 months.

An overall score was then calculated.

#### **Variables related to behaviours impacting on oral health**

- Child reported frequency of tooth-brushing. Data was divided into two groups; those who said that they brushed the recommended two times daily and those who said that they brushed once daily or less.
- Child reported usual dental attendance. Data was divided into two groups; those who attended regularly for dental examinations and those who only attended when they had “trouble with their teeth” or never attended.
- Child reported consumption of sugary food, grouped into two groups; four or more times in a day and less than four times a day.
- Child reported consumption of sugary drinks, grouped into two groups; four or more times in a day and less than four times a day.
- Self -reported orthodontic treatment. Previous research has highlighted that children who have experienced orthodontic treatment tend to be less anxious (12). Children were asked to report if they were having orthodontic treatment, if they have previously had orthodontic treatment or if teeth had been extracted for orthodontic reasons previously.

## **Results**

Six hundred and one 12 and 15 year olds were classified as dentally phobic (12.7%) with 4144 children classified as non phobic (87.3%). Mean 12.43 Std D 4.9.

The socio-demographic variables are shown in Table 1. Girls were more likely to be classed as dentally phobic. Eligibility for free school meals, used as an indicator of socio-economic status, indicated that there was no association with dental phobia.

**Table 1. Social and demographic features of participants**

The percentages stated refer to proportion of participants in either the non phobic or phobic group. For example, 51% of non phobic children were male and 49% were female. In the phobic group 30% were male and 70% were female.

		12 year and 15 year age group Non phobic	12 year and 15 year age group phobic	X <sup>2</sup>	P value
Gender	Male	2107 (51%)	181 (30%)	90.3	<0.002
	Female	2037 (49%)	420 (70%)		
Free school dinner eligibility	Yes	938 (24%)	149 (27%)	1.8	0.179
	No	2970 (76%)	411 (73%)		

The variables found during the clinical examination are shown in table 2. Phobic patients were more likely to have (i) active decay, (ii) teeth extracted due to decay and (iii) a higher PUFA score.

**Table 2 Clinical findings**

Variable		12 year and 15 year age group Non phobic	12 year and 15 year age group phobic	X <sup>2</sup>	P value

Number of teeth with active caries	0 teeth	2959 (71%)	375 (62%)	20.4	<0.002
	1+	1185 (29%)	226 (38%)		
Number of permanent teeth filled	0 teeth	2655 (64%)	377 (63%)	0.4	0.523
	1 + teeth	1489 (36%)	224 (37%)		
Number of primary teeth filled	0 teeth	4051 (98%)	587 (98%)	0.2	0.895
	1 + teeth	93 (2%)	14 (2%)		
Number of teeth extracted due to caries	0	3853 (93%)	536 (89%)	10.9	<0.002
	1+	291 (7%)	65 (11%)		
PUFA	Yes	106 (3%)	33 (6%)	16.2	<0.002
	No	4016 (97%)	561 (94%)		
Any decay experience	Yes	2207 (53%)	242 (40%)	8.9	0.03
	No	1937 (47%)	359 (60%)		

The reported oral health status of phobic versus non phobic groups is reported in Table 3.

Phobic children were more likely to have lower self rated dental and general health, and self reported bad breath.

**Table 3 -Child's self-reported oral health status**

Variable		12 year and 15 year age group Non phobic	12 year and 15 year age group phobic	$\chi^2$	P value
Self rated dental health	Very good/good	2947 (71%)	351 (59%)	38.6	<0.002
	Fair/worse	1175 (29%)	244 (41%)		
Self rate general health	Very good/good	3599 (87%)	471 (79%)	28.7	<0.002
	Fair/worse	526 (23%)	124 (31%)		
Reported toothache in the last 3 months	Not mentioned	3467 (84%)	482 (80%)	4.2	0.04
	Mentioned	676 (16%)	118 (20%)		
Reported sensitive tooth in the last 3 months	Not mentioned	2789 (67%)	381 (64%)	3.3	0.071
	Mentioned	1347 (33%)	217 (36%)		
Reported swollen or bleeding gums in the last 3 months	Not mentioned	3416 (86%)	470 (78%)	6.2	0.013
	Mentioned	724 (14%)	130 (22%)		
Reported broken tooth in the last 3 months	Not mentioned	3848 (93%)	545 (91%)	3.3	0.070
	Mentioned	294	55		



		(7%)	(9%)		
Reported mouth ulcers in the last 3 months	Not mentioned	3306 (80%)	473 (79%)	0.4	0.519
	Mentioned	835 (20%)	128 (21%)		
Reported bad breath in the last 3 months	Not mentioned	3403 (82%)	459 (76%)	12.1	<0.002
	Mentioned	743 (18%)	142 (24%)		

Regarding the variables relating to quality of life, the phobic children reported more difficulty eating, cleaning teeth, and reported that they felt different and had difficulty smiling, laughing and showing teeth without embarrassment as well as difficulty enjoying being with people. A sum of the scores given for the questions related to oral health affecting quality of life (Child OIDP) were analysed with Student's t-Test and showed a significant difference in means between the phobic and non phobic groups (p 0.001).

The variables relating to oral health related behaviour are shown in Table 4. Phobic children were more likely to brush less than twice a day and attend when only in trouble or not at all. There was no demonstrable relationship between dental phobia and sugar consumption.

**Table 4 – Oral health related behaviour**

Variable		12 year and 15 year age group Non phobic	12 year and 15 year age group phobic	$\chi^2$	P value
Frequency of brushing teeth	X2 a day	3218 (78%)	413 (70%)	21.1	<0.002
	X1 a day or less	889 (22%)	178 (30%)		
Pattern of attendance at dentist	For a check up	3439 (84%)	445 (74%)	33.9	<0.002
	Only when trouble/never	669 (16%)	156 (26%)		

Frequency of eating sugary food	4 times or more a day	2624 (64%)	376 (63%)	0.2	0.631
	Less than 4 times a day	1463 (36%)	219 (37%)		
Frequency of having sugary drink	4 times or more a day	1296 (32%)	194 (34%)	0.5	0.469
	Less than 4 times a day	2711 (68%)	379 (66%)		
Undergoing orthodontic treatment at time of survey	Yes	586 (14%)	68 (11%)	3.585	0.058
	No	3544 (86%)	532 (89%)		
Appliance worn in past	Yes	359 (10%)	40 (8%)	3.325	0.68
	No	3122 (90%)	477 (12%)		
(D) Any teeth extracted for orthodontic reasons?	No	3718 (90%)	543 (9%)	0.227	0.634
	Yes	426 (10%)	58 (91%)		

## **Discussion.**

The aim of this study was to assess the relationship of dental phobia and variables related to oral health. This study is among the first to assess this and utilized the data gathered from the well-planned Child Dental Health Survey. However, the use of MDAS as an anxiety measure may not have been ideal. The threshold value used to determine extreme dental anxiety has been validated on adult populations and may not be appropriate for use on the adolescent population group studied here.<sup>(30)</sup> However, some authors suggest there is no definitive anxiety measure for use with children.<sup>(29)</sup>

The overall picture described by the data is that dentally phobic children in these age groups are more likely to have dental related disease. In addition, this data clearly suggests that the child's oral health impacts on their quality of life, with phobic children reported that they were more likely to

have physical (eating, cleaning their teeth) and psychological difficulties (feeling different, embarrassed).

The prevalence of dental phobia in this population group is in line with the previous research. The relationship of dental anxiety and gender has been noted before,<sup>(20)</sup> and previous research has highlighted females are more likely to be dentally phobic.<sup>(3,7)</sup>

There was no significant difference between the phobic and non-phobic children in relation to treated caries such as primary or permanent teeth filled. There was a difference between the DMFT of non-phobic versus phobic patients, but this was not deemed as significant ( $p=0.03$ ). There was no demonstrable relationship between orthodontic treatment and dental phobia.

This data clearly suggests that there is an impact of the child's oral health on their quality of life. Phobic children reported that they were more likely to have physical (eating, cleaning their teeth) and psychological difficulties (feeling different, embarrassed) due to their oral health. This is in keeping with other studies findings on adults.<sup>(5,21)</sup> Self-esteem generally decreases during adolescence,<sup>(22)</sup> but the impact of poor oral health on a child's self-esteem may lead to a greater effect on development overtime. Further work needs to be undertaken to assess this.

Previous research has suggested that dentally phobic adults actively undertake preventative measures by using mouthwash.<sup>(7)</sup> This does not seem to apply to these children, but children are less likely to use mouthwash compared to adults; **indeed, in the UK children would not be able to purchase mouthwashes containing alcohol without assistance from a caregiver.**

There is a demonstrable link between caries experience and dental phobia in this and other research.<sup>(12)</sup> It can be hypothesised that phobic children experience a higher rate of dental disease due to three possible factors.

Firstly, the differences are due to a third factor related to phobia and oral health. For example, social class, which previous studies have shown a relationship between poorer oral health and an increased chance of dental phobia. However, this study showed no significant relationship between social class and phobia.

Secondly, there are differences in treatment approaches for phobic and non-phobic children. For example, dental phobics are more likely to choose a tooth extraction over extensive oral rehabilitation. This data may support this hypothesis with phobic children being more likely to have teeth extracted due to decay but no more likely to have teeth restored despite having more active decay or signs of untreated disease. Dentists may feel that there are fewer restorative options for phobic children, due to behavioural management problems. Recent research suggests that the level of behavioural knowledge of the profession is poor.<sup>(23)</sup> Improving the application of non pharmacological behavioural management may allow more conservative treatment on this phobic population group.

Finally, the differences in attendance are a result of fear. The link between irregular attendance and dental phobia is well supported and it has been postulated that this is due to escape/avoidance behaviour leading to the patient being negatively reinforced, resulting in greater avoidance of the dentist in the future.<sup>(24,25)</sup> However, with this age range the parents or caregiver are responsible for the child's attendance. We therefore must consider why the parent does not bring a child in need of treatment to the dentist. This may well be due to the parent not prioritising the child's oral health, a lack of knowledge about the importance of regular check ups, or the parent avoiding the dental environment due to their own dental fear or fear of the dentist's criticism. Sadly, no data was

available to analyse how the parent's anxiety affected the findings in this study. Further research is needed to fully investigate the reasons parents do not bring children to the dentist.

This research helps to provide a greater understanding of the oral health of dental phobics and demonstrates the need for the profession to investigate more fully the challenges faced by this population group.

However, the amount of theories postulated above demonstrates that this simple descriptive analysis has notable limitations. Therefore, there is a need to perform a multivariate analysis to fully understand the true relationships that we have touched upon here.

It is our opinion that this research demonstrates this population group face significant challenges due to dental anxiety, poor oral health and the significant impact this has on their self-esteem. Given the increasing evidence that behavioural therapies can lead to a long-term change in a person's beliefs about dentistry this seems to be a sizeable tragedy <sup>(26,27,28)</sup> These interventions can lead to a long term benefit for both the patient's health and the future economic burden on both the individual and the state.

## **References**

- (1) Poulton R, Waldie K.E., Thomson W.M., Locker D., Determinants of early- vs late-onset dental fear in a longitudinal-epidemiological study, *Behav Res Ther*, 2001, **39**, (7): 777-785.
- (2) O'Callaghan PM, Allen KD, Powell S. The efficacy of noncontingent escape for decreasing children's disruptive behavior during restorative dental treatment, *J Appl Behav Anal*, 2006, **39**: 161–171
- (3) Klingberg G, Broberg AG. Dental fear/anxiety and dental behaviour management problems in children and adolescents: a review of prevalence and concomitant psychological factors. *Int J Paediatr Dent*, 2007; **17**; 391-406
- (4) Edition F. Diagnostic and statistical manual of mental disorders. American Psychiatric Publishing, Arlington, VA; 2013.
- (5) Cohen SM, Fiske J, Newton JT. Behavioural dentistry: The impact of dental anxiety on daily living. *Br Dent J*, 2000, **14**;189(7):385-390.
- (6) McGrath C, Bedi R. The association between dental anxiety and oral health-related quality of life in Britain. *Community Dent Oral Epidemiol*. 2004, **32**(1):67-72.
- (7) Heidari E., Banerjee A, Newton J.T. Oral health status of non-phobic and dentally phobic individuals; a secondary analysis of the 2009 Adult Dental Health Survey. *Br Dent J*, 2015, **219**; 1-7
- (8) Abrahamsson K, Berggren U, Carlsson SG. Psychosocial aspects of dental and general fears in dental phobic patients. *Acta Odontol Scand*. 2000, **58**(1):37-43.
- (9) Hakeberg M, Berggren U. Changes in sick leave among Swedish dental patients after treatment for dental fear. *Community Dent Health*. 1993; **10**(1):23-29.
- (10) Salem K, Kousha M, Anissian A. et al. Dental Fear and Concomitant Factors in 3-6 Year-old Children. *J Dent Res Dent Clin Dent Prospects*, 2012, **6**(2):70-74.

- (11) Morgan, A.G., Rodd, D.R., Porritt, J.M., Baker, S.R., Creswell, C., Newton, J.T., Williams, C., Marshman, Z. Children's experience of dental anxiety, *Int J Paediatr Dent*, 2016, **27** (2) 87-97
- (12) Luoto A, Lahti S, Nevanperä T, Tolvanen M, Locker D. Oral-health-related quality of life among children with and without dental fear. *Int J Paediatr Dent*. 2009; **19**(2):115-120.
- (13) Children's Dental Health Survey Report 1: Attitudes, Behaviours and Children's Dental Health England, Wales and Northern Ireland, published 19 March 2015 retrieved 26/04/2018 from <https://files.digital.nhs.uk/publicationimport/pub17xxx/pub17137/cdhs2013-report1-attitudes-and-behaviours.pdf>
- (14) Humphris GM, Morrison T, Lindsay SJ. The Modified Dental Anxiety Scale: validation and United Kingdom norms. *Community Dent Health*. 1995 Sep. **12**(3):143-150.
- (15) Humphris G.M., Dyer T.A., Robinson P.G. The modified dental anxiety scale: UK general public population norms in 2008 with further psychometrics and effects of age *BMC Oral Health*, 2009, **9**:20
- (16) Humphris G, Crawford JR, Hill K, Gilbert A, Freeman R. UK population norms for the modified dental anxiety scale with percentile calculator: adult dental health survey 2009 results. *BMC Oral health*. 2013 Dec; **13**(1):29.
- (17) Porritt, J., Buchanan, H., Hall, M., Gilchrist, F. and Marshman, Z.. Assessing children's dental anxiety: a systematic review of current measures. *Community Dent Oral Epidemiol*, 2013 **41**(2).130-142.
- (18) Pitts NB, Chestnutt IG, Evans D, White D, Chadwick B, Steele JG. 1 Verifiable CPD Paper: The dentinal caries experience of children in the United Kingdom, 2003. *B Dent J*. 2006, **200**(6):313.
- (19) Gherunpong, S., Tsakos, G. and Sheiham, A. Developing and evaluating an oral health-related quality of life index for children; the CHILD-OIDP. *Community Dent Health*, 2004, **21**(2), 161-169.
- (20) Oosterink F, De Jongh A, Hoogstraten J. Prevalence of dental fear and phobia relative to other fear and phobia subtypes. *Eur J Oral Sci*. 2009. **117**(2):135-143.
- (21) Locker D, Jokovic A, Stephens M, Kenny D, Tompson B, Guyatt G. Family impact of child oral and oro-facial conditions. *Community Dent Oral Epidemiol*. 2002, **30**(6):438-448.
- (22) Robins RW, Trzesniewski KH. Self-esteem development across the lifespan. *Curr. Dir. Psychol. Sci*. 2005. **14**(3):158-162.
- (23) Coxon J, Hosey MT, Newton JT. Knowledge of behavioural management principles amongst specialist paediatric dental practitioners in the United Kingdom. *Behav Cogn Psychother*. 2017; **45**(2):185-192
- (24) Berggren U, Meynert G. Dental fear and avoidance: causes, symptoms, and consequences. *J. Am. Dent. Assoc*. 1984; **109**(2):247-251.
- (25) Thomson WM, Stewart JF, Carter KD, Spencer AJ. Dental anxiety among Australians. *Int Dent J*. 1996, **46**(4):320-324.
- (26) Kvale G, Berggren U, Milgrom P. Dental fear in adults: a meta-analysis of behavioural interventions. *Community Dent Oral Epidemiol* 2004; **32**: 250–264.

- (27) Shahnava S, Hedman E, Grindefjord M, Reuterskiöld L, Dahllöf G. Cognitive Behavioral Therapy for Children with Dental Anxiety: A Randomized Controlled Trial. *JDR Clin Trans Res*. 2016;**1**(3):234-243.
- (28) Kani E, Asimakopoulou K, Daly B, Hare J, Lewis J, Scambler S, Scott S, Newton JT. Characteristics of patients attending for cognitive behavioural therapy at one UK specialist unit for dental phobia and outcomes of treatment. *Br Dent J*. 2015 ;**219**(10):501-506.
- (29) Al-Namankany A, De Souza M, Ashley P. Evidence-based dentistry: analysis of dental anxiety scales for children. *Br Dent J*. 2012. **212**(5):219-222
- (30) Porritt J, Buchanan H, Hall M, Gilchrist F, Marshman Z. Assessing children's dental anxiety: a systematic review of current measures. *Community Dent Oral Epidemiol*. 2013. **41**(2):130-142.